



Focused Gastrointestinal Assessment

This course has been awarded one (1.0) contact hour.

First Published: October 4, 2004

Revised: September, 2018

Course Expires: September 30, 2021

Copyright © 2004 by AMN Healthcare in association with Interact Medical.

All Rights Reserved. Reproduction and distribution of these materials are prohibited without the express written authorization of AMN Healthcare.

Acknowledgements

RN.com acknowledges the valuable contributions of...

.....Kim Maryniak, PhD, RNC-NIC, NEA-BC

Conflict of Interest and Commercial Support

RN.com strives to present content in a fair and unbiased manner at all times, and has a full and fair disclosure policy that requires course faculty to declare any real or apparent commercial affiliation related to the content of this presentation. Note: Conflict of Interest is defined by ANCC as a situation in which an individual has an opportunity to affect educational content about products or services of a commercial interest with which he/she has a financial relationship.

The author of this course does not have any conflict of interest to declare.

The planners of the educational activity have no conflicts of interest to disclose.

There is no commercial support being used for this course.

Purpose and Objectives

The purpose of this course is to offer the healthcare provider an overview of basic gastrointestinal (GI) assessment including normal and abnormal findings.

After successful completion of this course, you will be able to:

1. Discuss the components of a focused gastrointestinal assessment.
2. Discuss history questions which will help focus your assessment.
3. Discuss specific assessment findings that are determined by the history and examination, including inspection, palpation, percussion, and auscultation.

Introduction

The functions of the gastrointestinal (GI) tract and its accessory organs are essential for life.

The process of digestion supplies nutrients to each and every cell in our body. If there is a disruption in any of these mechanisms, the whole body suffers.

This course will discuss specific information needed to obtain an adult patient's gastrointestinal history and will introduce exam techniques for your adult patient.

Physical exam techniques such as inspection, palpation, percussion, and auscultation will be highlighted.

Throughout the course, you will learn that deviations in your assessment findings could indicate potential gastrointestinal problems.

Glossary

Ascites - An abnormal accumulation of serous fluid in the abdominal cavity containing large amounts of protein and electrolytes.

Bulge - A protruding part; an outward curve or swelling.

Cirrhosis - Cirrhosis of the liver is a chronic disease of the liver characterized by the replacement of normal tissue with fibrous tissue and the loss of functional liver cells.

Digestion - The process by which food is converted into substances that can be absorbed and assimilated by the body.

Dysphagia - Difficulty in swallowing.

Esophageal varices - Abnormally dilated or swollen vessels in the esophagus, which can lead to bleeding.

Food allergy - An abnormally high sensitivity to certain foods.

Food intolerance - Inability to completely digest a type of food, usually due to an enzyme deficiency.

Hernia - The protrusion of an organ or other bodily structure through the wall that normally contains it; a rupture.

Mass - An aggregate of cells clumped together, such as a tumor.

Referred pain - Pain sensation experienced in one part of the body that is different to the actual area of pathology.

Spider nevi (or angioma) - A dilation of superficial capillaries with a central red dot from which blood vessels radiate.

Visceral pain - Pain related to the internal organs.

(Mosby, 2016; Venes, 2017)

Focused Gastrointestinal Assessment

When conducting a focused gastrointestinal assessment on your patient, both subjective and objective data are needed.

Components may include:

- Chief complaint
- Present health status
- Past health history
- Current lifestyle
- Psychosocial status
- Family history
- Physical assessment

Communication during the history and physical must be respectful and performed in a culturally-sensitive manner. Privacy is vital, and the healthcare professional needs to be aware of posture, body language, and tone of voice while interviewing the patient (Jarvis, 2016; Wilson & Giddens, 2017). Take into consideration that a patient's ethnicity and culture may affect the history that the patient provides.

Taking a Focused Gastrointestinal History

It is important to begin by obtaining a thorough history of abdominal or gastrointestinal complaints. You will need to elicit information about any complaints of gastrointestinal disease or disorders.

Gastrointestinal disease usually manifests as the presence of one or more of the following:

- Change in appetite
- Weight gain or loss
- Dysphagia
- Intolerance to certain foods
- Nausea and vomiting
- Change in bowel habits
- Abdominal pain

(Jarvis, 2016)

Appetite

Ask your patients if they have had any changes in appetite or food intake. If they have, ask for more information about the change. Appetite and eating can be influenced by many factors that may indicate gastrointestinal disease or that can be attributed to socioeconomic considerations such as food availability, family norms, peers, and cultural practices. A loss of taste sensation can contribute to loss of appetite and potentially result in poor nutrition, especially in older individuals. Attempts at voluntary control can be factors, such as dieting or eating disorders (National Institute of Mental Health [NIMH], 2016).

Weight Loss or Gain

Document any change in weight. If weight loss or gain is substantial or has happened rapidly, investigate further. Dieting to a body weight leaner than recommended health standards tends to be highly promoted by current fashion trends, sales campaigns for special foods, and is encouraged in some activities and professions. Young women are especially at risk for diet related alterations in normal gastrointestinal functions. Weight loss may also be associated with illness, while weight gain may be attributed to fluid retention or a mass (Jarvis, 2016).

Dysphagia

People with dysphagia have difficulty swallowing and may also experience pain while swallowing. Some people may be completely unable to swallow or may have trouble swallowing liquids, foods, or saliva. Eating becomes a challenge, making it difficult to take in enough calories and fluids to nourish the body.

Ask your patient if they have any difficulty swallowing and when the difficulty first occurred. More than 50 pairs of muscles and many nerves work to move food from the mouth to the stomach. It is important to note what the patient has difficulty swallowing (e.g. solids versus liquids), and the area that the patient feels is where food gets “stuck” (Porter & Kaplan, 2016).

People with diseases of the nervous system, such as cerebral palsy or Parkinson's disease, often have problems swallowing. Additionally, stroke or head injury may affect the coordination of the swallowing muscles or limit sensation in the mouth and throat. An infection or irritation can cause narrowing of the esophagus. People born with abnormalities of the swallowing mechanism may not be able to swallow normally. In addition, cancer of the head, neck, or esophagus may cause swallowing problems. Sometimes the treatment for these types of cancers can cause dysphagia. Injuries of the head, neck, and chest may also create swallowing problems (National Institute of Health [NIH], 2017).

Intolerance to Food

Ask your patient if they have any intolerance to certain foods. If so, ask which foods and the type of reaction to the food. Food intolerance should not be confused with food allergies. An intolerance to certain foods is generally based on the presence of a gastrointestinal imbalance such as having too little of a particular enzyme that can hinder proper breakdown and use of the food by the body. Food intolerance may be related to disorders such as celiac disease, insulin-dependent diabetes, and inflammatory bowel disease, or as a result of the aging process. Symptoms of intolerance to a particular food might include stomach discomfort, gas, bloating, burping, flatulence, abdominal pain, and diarrhea (NIH, 2011).

Nausea and Vomiting

Nausea and vomiting can be side effects of medications, a manifestation of many diseases, and can occur frequently in early pregnancy. Ask your patients about the frequency of these symptoms. Nausea and vomiting may also indicate food poisoning. Questions about types of food eaten in the past 24 hours should be asked to rule out potential poisoning.

If vomiting is present, you will want to ask about the amount, frequency, color, and odor of the vomitus. Ask if there is any blood in the vomit or if the vomit appears to be like coffee grounds. Hematemesis, or blood in the vomitus, is a common symptom of gastric or duodenal ulcers and may also indicate esophageal varices. Coffee ground emesis indicates an "old" gastrointestinal bleed. The old, partially digested blood appears to look like coffee grounds (Jarvis, 2016).

Changes in Bowel Habits

Particular emphasis should be placed on changes in bowel habits, as it is a common manifestation of gastrointestinal disease. The frequency, color, and consistency of bowel movements should be assessed. Assess the use of laxatives at this time.

Black, tarry stools may indicate an upper gastrointestinal bleed or may simply be from the ingestion of iron supplements or over the counter medications for gastrointestinal upset (Wilson & Giddens, 2017).

Bright red blood in the stools may indicate hemorrhoids or localized lower gastrointestinal bleeding.

Currant jelly stools are usually foul smelling and resemble maroon or purple colored jelly. The presence of currant jelly stools often indicates a massive bleeding episode and the patient's hemodynamic status must be assessed quickly (Wilson & Giddens, 2017).

Test Yourself

What can occur as a result of the aging process?

- A. Dysphagia
- B. Blood in the stools
- C. Increase in food intolerance

The correct answer is: C.

Past Gastrointestinal Disease and Medication History

Past Gastrointestinal Disease

Ask about any past history of gastrointestinal disorders such as ulcers, gall bladder disease, hepatitis, appendicitis, hernias. Ask the patient if they received treatment and if the treatment was successful. History should also include past abdominal surgeries, any abdominal problems after the surgery, and abdominal x-rays or tests (including colonoscopy) and their results (Jarvis, 2016).

Medication History

Many medications can produce gastrointestinal symptoms. Almost every class of drugs has the potential for gastrointestinal side effects. Most of the side effects include nausea, vomiting, diarrhea, and/or constipation.

Aspirin and non-steroidal anti-inflammatory drugs (NSAIDs) may cause abdominal pain and may increase the likelihood of gastrointestinal bleeding. Dietary supplements and the use of over the counter medications should also be included (Jarvis, 2016).

Social History and Lifestyle Risk Factors

In taking a complete history, it is important to address lifestyle risk factors and social behaviors that may contribute to unhealthy lifestyles and increase the risk of gastrointestinal disorders.

Ask your patients about the frequency and duration of alcohol consumption, caffeine intake, and cigarette smoking at this time. Alcohol can cause liver cirrhosis and esophageal varices. Cigarette smoking and regular ingestion of caffeine can lead to gastric reflux and gastric ulcers.

Also ask about recreational drug use such as marijuana, opiates, or amphetamines. The use of illicit drugs can increase or suppress appetite and affect GI function (Wilson & Giddens, 2017).

Test Yourself

Alcohol can cause liver cirrhosis and _____.

The correct response is esophageal varices.

Nutritional Assessment

Assessing nutritional status of your patients is important for several reasons. A thorough nutritional assessment will identify individuals at risk for malnutrition and provide baseline information for nutritional assessments in the future.

Some of your patients that will require a thorough nutritional assessment include those patients with:

- Recent unintentional weight loss
- Chemotherapy or radiation
- Recent weight gain
- Food allergies or intolerance
- Decreased appetite
- Multiple medications
- Alterations in sense of taste
- Dieting history
- Difficulty chewing or swallowing
- Vomiting
- Mobility problems
- Diarrhea
- Inability to feed self
- Recent surgery or major illness or injury
- Substance abuse
- Chronic conditions

- Potential for social isolation
- Low income

(Jarvis, 2016; Wilson & Giddens, 2017)

The Physical Exam

When performing a focused assessment, you will use at least one of the following four basic techniques during your physical exam: inspection, auscultation, percussion, and palpation. These techniques should be used in an organized manner from least disturbing or invasive to most invasive to the patient (Jarvis, 2016). Inspection is first, as it is non-invasive. Auscultation is performed following inspection; the abdomen should be auscultated before percussion or palpation to prevent production of false bowel sounds.

For accurate assessment of the abdomen, patient relaxation is essential. The patient should be comfortable with knees supported and arms at the sides, and should have an empty bladder. The environment should include a comfortable temperature, with good light.

The Physical Exam: Inspection

Visualization of the entire abdomen is needed. When assessing the abdomen, it is important to document the location of the physical exam finding. The abdomen can be divided into four or nine quadrants.

The Physical Exam: Inspection

With your patient in the supine position, inspect for:

- Bulges
- Masses
- Hernias
- Ascites
- Spider nevi
- Enlarged veins
- Pulsations or movements
- Inability to lie flat

Normally, blood vessels are not evident on the abdomen. However they may be present in the elderly or pregnant client due to the loss of subcutaneous fat.

During inspection ask your patient to lift their head slightly. If you notice a protrusion around the umbilicus or any incisions, a hernia may be present (Jarvis, 2016).

The Physical Exam: Auscultation

You should always auscultate the abdomen after inspection and before percussion or palpation so you do not produce false bowel sounds by percussion or palpation. Auscultation should begin in the right lower quadrant. If bowel sounds are not heard, in order to determine if bowel sounds are truly absent, listen for a total of five minutes (Jarvis, 2016).

Bowel sounds echo the underlying movements of the intestines. It is normal to hear high-pitched clicking and gurgling sounds approximately every five to 15 seconds.

It is suggested that you listen to bowel sounds for a full minute before determining if they are normal, hypoactive, or hyperactive. Refer to the table to see how different bowel sounds are produced and what they may indicate.

An example of a video demonstrating abdominal auscultation can be viewed at:

<https://www.youtube.com/watch?v=7VG0rx5nkGw>

Table of Bowel Sounds

Bowel Sound	How it is Produced	Possible Cause
Normal Bowel Sounds	Intestines transporting fluid and digested food through intestinal lumen at normal rate. Sounds are approximately every 5 to 15 seconds.	<ul style="list-style-type: none"> • Normally functioning intestine
Hypoactive Bowel Sounds	Intestines transporting fluid and digested food through intestinal lumen at a decreased rate probably due to inactivity of smooth muscle in the bowel. Sounds are approximately every 20 to 30 seconds; can be longer.	<ul style="list-style-type: none"> • Paralytic ileus • Peritonitis • Decreased bowel motility • Late intestinal obstruction
Hyperactive Bowel Sounds	Intestines transporting fluid and digested food through intestinal lumen at an increased rate probably due to rapid passage of air and fluid through the intestines. Sounds can be as frequent as every second.	<ul style="list-style-type: none"> • Diarrhea • Early intestinal obstruction • Gastroenteritis • Anxiety
High-pitched Rushing or Tinkling Sounds (Borborvomi)	<u>Hyperperistalsis</u> from intestinal straining to push fluid and/or air past an obstruction, or fluid and/or air under pressure. Very loud sounds; may be heard without a stethoscope.	<ul style="list-style-type: none"> • Intestinal obstruction • Dilated bowel loops • Fecal impaction • Gastroenteritis
Absent Bowel Sounds	Absence of intestinal motility Ominous finding	<ul style="list-style-type: none"> • Peritonitis • Late obstruction (ileus) • Perforation • Trauma
Abdominal Bruits	Whooshing sound over an artery from increased turbulence of blood flow in that artery	<ul style="list-style-type: none"> • Aneurysm • Thin, emaciated patient • Renal artery stenosis

(Jarvis, 2011; Shaw 2012)

The Physical Exam: Percussion

Percussion is used to elicit tenderness or sounds that give clues to underlying problems. When percussing directly over suspected areas of tenderness, monitor the patient for signs of discomfort. Percussion requires skill and practice, by pressing the distal part of the non-dominant hand's middle finger firmly on the patient's body part. The rest of the hand should be off the body surface. Using the middle finger of the dominant hand, tapping directly over the point where the other finger connects with the patient's skin. Keep the fingers perpendicular. Listen to the sounds produced by the percussion.

The Physical Exam: Percussion

When examining the abdomen, percuss for general tympany, liver span, and splenic dullness. Tympany should be the predominant sound when percussing the abdomen. Air "floats" to the top of the abdomen in the supine position and tympany reflects a drum-like sound (Wilson & Giddens, 2017).

Dullness is usually heard over solid organs or masses such as the liver, spleen, or a full bladder (Wilson & Giddens, 2017).

The Physical Exam: Percussion

Percussing over the kidneys does not usually produce pain or discomfort. If tenderness is present, a urinary tract infection or kidney inflammation may be present.

Costovertebral angle tenderness may be elicited when the patient is in a standing or upright position. Place the palm of your non-dominant hand near the posterior costovertebral margin over the kidney. Gently, but firmly, tap on your hand with the fist of your other hand. An example of a video demonstrating abdominal percussion can be viewed at: <http://www.youtube.com/watch?v=5ERuM1JDYAA>

To determine if abdominal distention is due to fluid or air, you may want to ask a nursing assistant or another nurse to assist you in percussing a fluid wave. When percussing a fluid wave, your assistant should place her arm and hand along the mid-line of the patient's abdomen, with the patient in the supine position. Her arm should be placed firmly on the abdomen to prevent the transmission of fat waves. You should then place your palm of one of your hands in the lateral lumbar region of the patient's abdomen. With your other hand, quickly pat or tap the other lateral lumbar region of your patient's abdomen. If a fluid wave is present, as with ascites, you will feel the resulting wave with your opposite hand. If the distention is due to air you will not feel any wave.

Did You Know?

Tympany should be the predominant sound when percussing the abdomen. Air "floats" to the top of the abdomen in the supine position and tympany reflects a drum-like sound (Jarvis, 2016).

The Physical Exam: Palpation

Palpation is another commonly used physical exam technique that requires you to touch your patient with different parts of your hand using different strength pressures. During light palpation, you press the skin about $\frac{1}{2}$ inch to $\frac{3}{4}$ inch with the pads of your fingers. When using deep palpation, use your finger pads and compress the skin about $1\frac{1}{2}$ to 2 inches. Palpate lightly then deeply noting any muscle guarding, rigidity, masses or tenderness. Palpate tender areas last. Only if indicated, palpate the liver margins, the spleen or the kidneys and percuss the abdomen for general tympany, liver span, splenic dullness, costovertebral angle tenderness, presence of fluid wave, or shifting dullness with ascites (Jarvis, 2016).

Palpation allows you to assess for texture, tenderness, temperature, moisture, pulsations, masses, and internal organs (Wilson & Giddens, 2017). Normally, you should elicit no tenderness on either light or deep palpation of the abdomen. If inguinal lymph nodes are palpated, they should be small and freely moveable.

Test Yourself

During light palpation compress the skin:

- A. $\frac{1}{2}$ inch to $\frac{3}{4}$ inch
- B. $\frac{1}{2}$ inch to 2 inches
- C. $1\frac{1}{2}$ inches to 2 inches
- D. $1\frac{1}{2}$ inches to 3 inches

The correct answer is: A.

Abdominal Pain

Introduction

If your patient is experiencing abdominal pain, have them point to the exact location of the pain.

Abdominal pain can be classified as:

- Visceral
- Parietal
- Referred

Visceral Pain

Visceral pain is usually described as dull, crampy, squeezing, or aching. It can be constant or intermittent. The pain may be difficult to localize and may be located over an abdominal organ (Jarvis, 2016).

Parietal Pain

Parietal pain is usually from inflammation over the peritoneum. Peritoneal inflammation usually indicates an underlying emergency and should be assessed quickly. Parietal pain is usually intense, constant, and on one side. It can be aggravated by extension of the lower extremity on the affected side, coughing, or eliciting rebound tenderness (Jarvis, 2016).

Referred Pain

Referred pain is usually visceral pain that is felt in another area of the body when a common nerve pathway is shared. It occurs with specific gastrointestinal disorders such as appendicitis (can cause umbilical pain in early stages), gall bladder disease (referred to right upper scapula), and pancreatitis (referred to the mid-back) (Jarvis, 2016).

Mnemonic for Pain Assessment

Introduction

In general, the mnemonic, PQRST, is very useful in assessing abdominal pain and other gastrointestinal symptoms, such as distention, nausea, and vomiting. It provides a methodology in which communication to other healthcare providers will be efficient and informative.

After eliciting information about any experienced signs or symptoms of gastrointestinal disease, ask about your patients past abdominal or gastrointestinal history, medications, and nutritional status.

P: Provocative or Palliative: What makes the pain or symptom(s) better or worse?

Q: Quality: Describe the pain or symptom(s) (burning, dull, sharp)

R: Region or Radiation: Where in the body does the pain or symptom(s) occur? Is there radiation or extension of the pain or symptom(s) to another area of the abdomen?

S: Severity: On a scale of 1-10, (10 being the worst) how bad is the pain or symptom(s)? Another visual pain scale may be appropriate for patients that are unable to identify with this scale.

T: Timing: Does it occur in association with something else? (e.g. eating, exertion, movement)

Assessing Abdominal Pain: Muscle Tests

The patient history is extremely important in assessing abdominal pain. Pain may be chronic or acute and related to inflammation, infection, allergy, or food intolerance. It can also result from trauma or obstruction. There are also a few physical exam techniques that can be used to assess acute abdominal pain. These are the iliopsoas muscle test, obturator test, and Blumberg test (Porter & Kaplan, 2016; Wilson & Giddens, 2017).

Iliopsoas Muscle Test

The iliopsoas muscle test is used most often when acute abdominal pain is present and appendicitis is suspected.

When your patient is lying in the supine position ask him or her to lift their right leg straight up, flexing only at the hip. Push down on the lower part of the thigh when your patient is trying to hold their leg up. If the patient feels pain in the iliopsoas muscle (the right lower quadrant of the abdomen) the test is positive and may indicate a perforated or inflamed appendix.

Anticipate further investigatory tests to confirm a suspected diagnosis (Porter & Kaplan, 2016).

The Obturator Test

The obturator muscle test is also performed when acute abdominal pain is present and appendicitis is suspected. When your patient is lying in the supine position ask him or her to lift their right leg straight up, flexing at the hip, and 90 degrees at the knee. Hold the ankle and rotate the leg internally and externally. If the patient feels pain in the area of the internal obturator muscle (the right lower quadrant of the abdomen and pelvis) the test is positive and may also indicate a perforated or inflamed appendix (Porter & Kaplan, 2016).

The Blumberg Sign

Blumberg Sign is also known as rebound abdominal tenderness. Choose a site away from the suspected area of tenderness. Holding your hand 90 degrees to the abdomen, press inward deeply, then release quickly. Pain on release of pressure is an indicator of peritoneal irritation (Porter & Kaplan, 2016).

Assessing and Interpreting Associated Laboratory Values

There are many common lab values that will help you in your assessment of your patient's gastrointestinal system and accessory organs. Lab values should be looked at collectively in the context of a complete abdominal history and examination. The following table illustrates examples of lab values and the possible related gastrointestinal disturbance.

Porter, & Kaplan, 2016

<i>*Normal lab value reference ranges differ between labs and institutions. Check with your facility for normal ranges.</i>			
Lab	Normal Value	Alteration	Potential Gastrointestinal Cause of Abnormal Value
Lipase	7-60 u/L	↑	Pancreatitis
Amylase	30-170 u/L	↑	Pancreatitis
Calcium	8.5-10.3 mg/dL	↓	Pancreatitis, malnutrition
Platelets	130-400 x 10 ³ /mm ³	↓	Liver dysfunction, cirrhosis, hepatitis, GI bleed
AST	< 42 u/L	↑	Liver dysfunction, cirrhosis, hepatitis
ALT	< 48 u/L	↑	Liver dysfunction, cirrhosis, hepatitis
Fibrinogen	200-400 mg/ dL	↓	Liver dysfunction, cirrhosis, hepatitis
Prothrombin Time (PT)	(PT) 10.0-12.5 sec	↑	Liver dysfunction, cirrhosis, hepatitis
Albumin	3.5-5.0 g/dL	↓	Liver dysfunction, cirrhosis, hepatitis, malnutrition
Bilirubin	≤ 1.3 mg/dL	↑	Liver dysfunction, cirrhosis, hepatitis, <u>cholecystitis</u>
Ammonia	0.17-0.80 mcg/mL	↑	Liver failure
Hemoglobin	12.0-17.2 g/dL	↓	GI bleed, hemorrhagic pancreatitis
Hematocrit	35-50%	↓	GI bleed, hemorrhagic pancreatitis
Electrolytes	variable	↑	<u>Hemoconcentration</u> in early GI bleed or hemorrhagic pancreatitis
BUN	7-30 mg/dL	↑	<u>Hemoconcentration</u> & absorption of protein (blood) in GI bleed hemorrhagic pancreatitis
WBC	3.8-10.8 x 10 ³ /mm ³	↑	Infection of stress response of pancreatitis, GI bleed

(Merck Manual Online, 2013)

Porter, & Kaplan, 2016

Conclusion

Digestion, transport, and absorption are the processes by which the digestive system supplies nutrients to each and every cell of our body. If there is a disruption to this process, the whole body suffers.

By asking specific questions about a patient's gastrointestinal history and performing focused abdominal exam techniques for your adult patient, you will be able to assess for the slightest changes in gastrointestinal function.

Alterations in your gastrointestinal assessment findings could indicate potential problems.

Being knowledgeable about the focused, gastrointestinal assessment will allow you to intervene quickly and appropriately for gastrointestinal disorders.

References

Jarvis, C. (2016). *Physical examination and health assessment* (7th ed.). St. Louis: W.B. Saunders.

National Institute of Health (NIH). (2017). Dysphagia. Retrieved from <http://www.nidcd.nih.gov/health/voice/pages/dysph.aspx>

National Institute of Mental Health (NIMH). (2016). Eating disorders. Retrieved from <https://www.nimh.nih.gov/health/topics/eating-disorders/index.shtml>

Mosby Company. (2016). *Mosby's medical dictionary* (10th ed.). New York: Elsevier.

Porter, R., & Kaplan, J. (eds.). (2016). *Merck manual professional version*. Whitehouse Station, NJ: Merck Sharp & Dohme Corp.

Venes, D. (ed.) (2017). *Tabers® cyclopedic medical dictionary* (23rd ed.). Philadelphia: F.A. Davis Co.

Wilson, S. & Giddens, J. (2017). *Health assessment for nursing practice* (6th ed.). St. Louis, MO: Elsevier.

At the time this course was constructed all URL's in the reference list were current and accessible. RN.com is committed to providing healthcare professionals with the most up-to-date information available.

© Copyright 2004, AMN Healthcare, Inc.

Disclaimer

This publication is intended solely for the educational use of healthcare professionals taking this course, for credit, from RN.com, in accordance with RN.com terms of use. It is designed to assist healthcare professionals, including nurses, in addressing many issues associated with healthcare. The guidance provided in this publication is general in nature, and is not designed to address any specific situation. As always, in assessing and responding to specific patient care situations, healthcare professionals must use their judgment, as well as follow the policies of their organization and any applicable law. This publication in no way absolves facilities of their responsibility for the appropriate orientation of healthcare professionals. Healthcare organizations using this publication as a part of their own orientation processes should review the contents of this publication to ensure accuracy and compliance before using this publication. Healthcare providers, hospitals and facilities that use this publication agree to defend and indemnify, and shall hold RN.com, including its parent(s), subsidiaries, affiliates, officers/directors, and employees from liability resulting from the use of this publication. The contents of this publication may not be reproduced without written permission from RN.com.

Participants are advised that the accredited status of RN.com does not imply endorsement by the provider or ANCC of any products/therapeutics mentioned in this

course. The information in the course is for educational purposes only. There is no “off label” usage of drugs or products discussed in this course.

You may find that both generic and trade names are used in courses produced by RN.com. The use of trade names does not indicate any preference of one trade named agent or company over another. Trade names are provided to enhance recognition of agents described in the course.

Note: All dosages given are for adults unless otherwise stated. The information on medications contained in this course is not meant to be prescriptive or all-encompassing. You are encouraged to consult with physicians and pharmacists about all medication issues for your patients.